

Field-induced criticality in a gapped quantum magnet with bond disorder.

D. Hübner,¹ S. Zhao,¹ M. Månsson,¹ T. Yankova,^{1,2} E. Ressouche,³
C. Niedermayer,⁴ M. Laver,^{4,5,6} S.N. Gvasaliya,¹ and A. Zheludev^{1,*}

¹*Neutron Scattering and Magnetism, Laboratory for Solid State Physics, ETH Zurich, Switzerland.*

²*Chemistry Dept., Lomonosov Moscow State University, Russia.*

³*INAC/SPSMS-MDN, CEA/Grenoble, 17 rue des Martyrs, 38054 Grenoble Cedex 9, France*

⁴*Laboratory for Neutron Scattering, Paul Scherrer Institute, CH-5232 Villigen, Switzerland.*

⁵*Materials Research Division, Ris DTU, Technical University of Denmark, DK-4000 Roskilde, Denmark.*

⁶*Nano Science Center, Niels Bohr Institute, University of Copenhagen, DK-2100 Copenhagen, Denmark.*

(Dated: March 14, 2012)

Neutron diffraction and calorimetric measurements are used to study the field-induced quantum phase transition in piperazinium- $\text{Cu}_2(\text{Cl}_{1-x}\text{Br}_x)_6$ ($x = 0$, $x = 3.5\%$ and $x = 7.5\%$), a prototypical quantum antiferromagnet with random bonds. The critical indexes ϕ and β are determined. The findings contradict some original predictions for Bose Glass based on the assumption $z = d$, but are consistent with recent theoretical results implying $z \leq d$. Inelastic neutron experiments reveal that disorder has a profound effect on the lowest-energy magnetic gap excitation in the system.

PACS numbers: 75.10.Jm, 64.70.Tg, 75.40.Cx, 72.15.Rn

The persistent interest in quantum magnets is driven by their use as prototypes for the study of quantum critical phenomena. In this context, much attention has been given to field-induced phase transitions in gapped spin systems with a singlet ground state, the so-called quantum spin liquids. Such transitions are described in terms of a Bose Einstein condensation (BEC) of magnons.¹ The specific advantage of magnetic realizations of BEC lies in the relatively easy experimental access to- and control of the key relevant quantities, notably the effective chemical potential, boson density and BEC order parameter. In magnetic BEC, the latter correspond to magnetic field, magnetization and antiferromagnetic order, respectively.

A particular area of recent interest is BEC in the presence of a spatially random potential. Such disorder leads to a localization of bosons and the appearance of a novel quantum phase, the so-called Bose Glass.²⁻⁴ Most importantly, it changes the universality class of the quantum critical point. While in the disorder-free case it is Mean Field (MF) like with dynamical critical exponent $z = 2$,¹ very unusual scaling with z equal to dimensionality ($z = d$) was predicted for disordered systems.³ This premise was recently challenged,^{5,6} suggesting that in fact $z \leq d$.^{7,8} Quantum magnetic materials offer a unique opportunity of addressing this important matter of debate experimentally. A random potential for magnons can be created by randomizing the strength of magnetic interactions. This is achieved through a spatially random chemical modification. Indications of the magnetic Bose Glass⁹⁻¹¹ and a modification of the field-induced ordering transition¹²⁻¹⁴ have been reported. However, experiments only added to the confusion regarding critical indexes and scaling. Recent studies of $\text{Tl}_{0.64}\text{K}_{0.36}\text{CuCl}_3$ seemed to validate the $z = d$ hypothesis.¹² Other experiments, on the organic compound DTN,¹³ yielded quite different critical exponents, and were supported by Density Matrix Renormalization Group (DMRG) calculations. In the present work we seek to resolve the

issue through neutron and calorimetric studies of the chemically disordered quantum magnets piperazinium- $\text{Cu}_2(\text{Cl}_{1-x}\text{Br}_x)_6$ (PHCX). We determine the critical exponents ϕ and β , the latter previously unmeasured for this type of transition. We find experimental values that are clearly inconsistent with predictions based on $z = d$.

One rationale for selecting PHCX as a test material is that the disorder-free Cl-rich ($x = 0$) compound, (PHCC) is an exceptionally well-characterized Heisenberg quantum magnet.¹⁵⁻¹⁸ It features a complex quasi two-dimensional spin network of $S = 1/2$ Cu^{2+} ions bridged by Cu-Cl-Cl-Cu superexchange pathways, see Fig.1 in Ref.15. The exact Hamiltonian for this system remains unknown with 6 to 8 possible superexchange pathways within the Cu_2Cl_6 layers. However, for the physics discussed here relevant ramification is the spin liquid ground state and the gapped ($\Delta = 1$ meV) magnon spectrum with bandwidth of about 1.7 meV. In magnetic fields exceeding $\mu_0 H_c = 7.5$ T, PHCC undergoes a quantum phase transition to a state with spontaneous antiferromagnetic (AF) order perpendicular to the direction of applied field. The critical properties of this transition have been investigated using a variety of techniques.¹⁷ The two critical exponents studied in this work are the order parameter exponent β and the so-called crossover exponent ϕ . The former defines the field dependence of the AF ordered moment at $T \rightarrow 0$: $|\langle \mathbf{S}_\perp \rangle| \sim (H - H_c)^\beta$, where $H_c \equiv H_c(T = 0)$. The crossover index ϕ defines the phase boundary of the ordered state on the $H - T$ phase diagram: $T_c(H) \sim (H - H_c)^\phi$ or $T_c(H)^{1/\phi} \sim (H - H_c)$.

Br substitution in PHCX is expected to locally affect the bond angles in the halogen-mediated superexchange routes, and thereby to realize the random bond model. This mechanism was previously exploited in the study of random-bond systems IPA-CuCl_3 ,^{9,11} $\text{Sul-Cu}_2\text{Cl}_4$ (Ref. 14) and DTN.¹³ High quality PHCX single crystals were grown from solution as described in

Ref. 19. Single crystal X-ray diffraction confirmed that for Br content x up to at least 10%, the crystal symmetry remains intact, space group ($P\bar{1}$). The lattice parameters are $a = 7.9691(5)\text{\AA}$, $b = 7.0348(5)\text{\AA}$, $c = 6.0836(4)\text{\AA}$, $\alpha = 111.083(3)^\circ$, $\beta = 99.947(3)^\circ$, $\gamma = 81.287(4)^\circ$ for the pure material with $x = 0$,²⁰ and the cell dimensions increase linearly with x . Here x stands for nominal Br concentration in the starting solution. Studies on similar compounds have shown a very good correspondence between nominal and final dopant concentrations.^{14,19} Most of the work reported here was done on perfectly faceted single crystals with $x = 3.5\%$ and $x = 7.5\%$, where $a = 7.9799(2)\text{\AA}$, $b = 7.0406(2)\text{\AA}$, $c = 6.0909(10)\text{\AA}$ and $a = 7.9905(3)\text{\AA}$, $b = 7.0454(2)\text{\AA}$, $c = 6.1000(2)\text{\AA}$, respectively, with the triclinic angles indistinguishable from the pure compound's.

The crossover exponent was derived from calorimetric measurements. For $x = 0$, $x = 3.5\%$ and $x = 7.5\%$, the data were collected using a 14 T Quantum Design PPMS with a ^3He - ^4He dilution insert, both as a function of temperature and magnetic field applied along the a^* axis. Typical scans are shown in Fig. 1a,b (symbols). Sharp lambda-anomalies are apparent in all cases and mark the ordering transition. The apparent broadening at low T for $x = 7.5\%$ sample is of instrumental origin. It is caused by differences in the slope of the phase boundary in conjunction with finite temperature resolution. In the constant-field scans, the actual transition temperatures T_c were determined by fitting the data at each field, in the vicinity of the peaks, to an empirical form of power law with broadening.²¹ A similar approach was used to obtain the temperature dependence of the critical field H_c from constant-temperature scans. In this case the fit function was a composite of two half-Lorentzians of non-equal widths and common maxima. The resulting phase boundaries are plotted in Fig. 2 (symbols). Clearly, the introduction of disorder shifts the transition to higher fields and lower temperatures.

Any more quantitative analysis has to take into account the experimental errors. For consistency, we only used the more comprehensive data set obtained in constant temperature scans (empty symbols in Fig. 2). The error bar on H_c was estimated as the standard deviation of this parameter in the least squares fits described above and is significantly smaller than the symbol size in the plot. The dominant error on T_c is of instrumental origin, and due to the use of the temperature relaxation measurement method. It is defined by the magnitude of applied heat pulse (3% of the starting temperature in our case).

The exponent ϕ is obtained by weighted least squares fit of the power law function to the experimental field-dependence of T_c accounting for experimental errors on both. This deceptively simple procedure is actually quite delicate. The fit is performed in a progressively shrinking data range $T < T_{\max} \rightarrow 0$. In Figs. 2b-2d, we show the fitted values of ϕ for the three materials plotted against $T_{\max}$. The range used for the final fits is to be selected




FIG. 1. (color online) Symbols: Heat capacity of PHCX with varying Br concentration x measured (a) as a function of magnetic field at $T=0.5\text{K}$ and (b) as a function of temperature at $H = 14\text{ T}$. The solid lines are fits to the data as described in the text.

	$\mu_0 H_c\text{ T}$	ϕ	β
PHCX 0.0% Br	7.67 ± 0.02	0.72 ± 0.04	0.40 ± 0.01 [17]
PHCX 3.5% Br	9.14 ± 0.08	1.02 ± 0.16	0.45 ± 0.05
PHCX 7.5% Br	10.82 ± 0.14	0.85 ± 0.10	0.42 ± 0.02
DTN 0.8% Br [13]	1.07 ± 0.01	1.1 ± 0.2	
Tl _{0.64} K _{0.36} CuCl ₃ [12]	2.7 ± 0.6	1.7	
$z = d$ scaling [3]		≥ 2	
QMC [7]		1.54	0.61
QMC [13]		1.06 ± 0.09	
BEC (MF)		0.667	0.5

TABLE I. Experimental estimates for the critical field H_c in $\mathbf{H}||\mathbf{a}^*$ orientation and the critical exponents ϕ (from calorimetric data) and β (from neutron diffraction) of the field-induced quantum phase transition in PHCX in comparison with recent measurements on similar systems and theoretical results, whose relevance is discussed in the text.

based on two conflicting requirements. It has to be as narrow as possible, in order to access the true critical region. This desire is counteracted by the divergence of confidence intervals at $T_{\max} \rightarrow 0$. As a rule of thumb, one can choose the *largest* $T_{\max}$ that gives a value of ϕ within the confidence intervals obtained using any narrower ranges. Further reducing the fitting range does not produce a statistically significant change in the result.

Returning to our case of PHCC, based on the plots in Fig. 2 and the above-mentioned rule of thumb, for the $x = 0$, $x = 3.5\%$ and $x = 7.5\%$ materials we have selected fitting ranges $T_{\max} = 1.75\text{ K}$, 0.9 K and 1.15 K , respectively. The resulting values of ϕ and H_c are summarized in Table I. Power laws derived from these parameter values are shown as solid lines up to $T = T_{\max}$ in Fig. 2.

The behavior of the magnetic BEC order parameter was studied in neutron diffraction experiments. For the




FIG. 2. (color online) (a) Phase boundaries of ordered state for Br-free, 3.5% and 7.5% Br substituted PHCX. Open and filled circles are from $C(H)$ and $C(T)$ measurements, respectively. Solid lines are a power laws $H - H_c \sim T_c(H)^{1/\phi}$ with critical fields H_c and crossover exponents ϕ derived from the data as described in the text. (b-d) Crossover exponent ϕ obtained from least squares fit as a function of fitting window given by $T < T_{max}$. Filled symbols show the value quoted in Table I satisfying our best fit criteria described in the text.

$x = 3.5\%$ and $x = 7.5\%$ samples these were carried out at the CRG-CEA D23 diffractometer at ILL and the RITA-II 3-axis spectrometer at PSI, respectively. In the two experiments, we used Cu monochromator to produce neutrons of incident energies 50 meV at ILL and pyrolytic graphite for 4.6 meV neutrons with Be filter in front of the analyzer at PSI. Sample environment was in all cases a split coil cryomagnet with a ^3He - ^4He dilution cryostat. The magnetic field was applied along a^* in the first and along the b crystallographic axis in the second experiment. As previously reported for PHCC,¹⁷ long range order in the BEC phase is marked by the appearance of new magnetic Bragg reflections half-integer reciprocal-lattice points $(h + 1/2, 0, l + 1/2)$. The inset in Fig.3b shows typical scans across the $(0.5, 0, -0.5)$ position measured in the $x = 7.5\%$ sample at $T = 0.5$ K in the BEC




FIG. 3. (color online) *Upper panels:* Neutron diffraction intensity of the $(\frac{1}{2}0\frac{1}{2})$ peak measured in PHCX, a) as a function of magnetic field at $T = 160$ mK, b) as a function of temperature. Open and solid symbols correspond to zero field cooling and field cooling, respectively. Squares (circles) correspond to 3.5% ($x = 7.5\%$) Br content. Lower left corner: the trajectory in (H, T) space for the measurements on the $x = 7.5\%$ material. Inset: rocking curve measured at $T = 0.5$ K (filled) and $T = 2$ K (empty circles) in that sample. *Lower panels:* order parameter exponents as obtained by shrinking the fitting interval analysis of the 3.5% (panel c) and 7.5% Br (panel d) samples data. Filled symbols show the best fit values of β .

phase, and the background collected at $T = 2$ K in the paramagnetic state. To within experimental error, all magnetic reflections are resolution-limited.

The critical index β was determined from measurements of the $(0.5, 0, -0.5)$ Bragg peak intensity in zero field cooled (ZFC) samples. These data were taken upon increasing the applied field at $T = 160$ mK and are plotted in Fig.3a. β was then determined in power law fits, assuming that peak intensity scales as the square of the order parameter. As for ϕ , the fits for β were performed for a series of shrinking fitting intervals, of magnetic field $\delta H = H - H_c$ in this case. The fit results and standard deviation confidence intervals are plotted against δH in Figs. 3c,d. Following the reasoning described earlier, in our final analysis we have selected $\mu_0 \delta H = 1.86$ T and 2.15 T, for $x = 3.5\%$ and $x = 7.5\%$, respectively. The resulting exponents are summarized in Table I. In order to properly characterize the quantum critical point, one has to know β in the limit $T \rightarrow 0$. Previous studies of PHCC suggest that β remains practically constant for




FIG. 4. (color online) Inelastic neutron scattering intensity at (0.5,0.5,-0.5) in energy transfer range from 0.5 to 2.5 meV for 3.5% (green squares) and $x = 7.5\%$ Br (red circles) content samples. Corresponding solid lines are the fitted Lorentzian energy profiles convoluted with the 4-dimensional instrumental resolution function. For 0.0% Br sample resolution convoluted simulation is shown for comparison using parameters from Ref.15

$T < 0.3$ K.¹⁷ We have additionally verified this behavior for our 3.5% PHCX sample. Thus, the 160 mK measurements are good estimates of the zero-temperature values. Comparing our results with those reported in Ref. 17, we conclude that disorder does not significantly change the criticality of the order parameter.

This is not to say that the effect of disorder is negligible. Inelastic neutron scattering experiments performed at zero field on the TASP 3-axis spectrometer²² at PSI using 3.5 meV incident-energy neutrons and Be filter reveal that while the magnons are sharp in PHCC,¹⁷ they acquire a considerable intrinsic width in disordered samples. Typical constant- q scans at the AF zone-center (0.5,0.5,-0.5) in the $x=3.5\%$ and $x=7.5\%$ samples are shown in Fig. 4 by the open symbols. The solid lines are fits of Lorentzian energy profiles convoluted with the 4-dimensional instrumental resolution function²³. For $x=0\%$ the line is a simulation assuming that peak shape is determined solely by resolution. The observed increase of gap energy with Br substitution is fully consistent

with the observed increase of critical field and may be attributed to a change in the disorder-averaged exchange constants. However, the large intrinsic energy width $\Gamma = 0.1$ meV and 0.25 meV for $x=3.5\%$ and $x=7.5\%$ samples, respectively, is entirely a disorder-induced effect²⁴.

Our present diffraction experiments show that disorder also affects the behavior of the order parameter deep within the high-field phase. In the 7.5% material the intensity of the magnetic reflection exhibits a history dependence. This is illustrated in Fig. 3a,b. Here we show data collected along a trajectory illustrated in the inset. No such behavior was reported for the disorder-free material, but a similar, and much stronger history effect was previously studied in a related system IPA-(Cu_{1-x}Br_x)Cl₃.¹¹ Related phase transition scenario that could lead to hysteretic behavior was discussed in Ref. 26.

Theoretical and numerical predictions for the two measured critical exponents are shown in Table I for a direct comparison with our experiments. In the absence of disorder, one expects MF values¹⁷ $\beta = 1/2$ and $\phi = 2/3$, in reasonable agreement with the data for PHCC. Assuming $z = d$ the disordered case gives much larger value for the crossover exponent $\phi \geq 2$.²⁵ This prediction is *clearly inconsistent with our experiments*. More recent numerical work^{7,8,13} suggested smaller values of the exponent. Those from Refs. 7 and 8 are less relevant to 3-dimensional ordering in PHCX, as they are obtained for a 2-dimensional model. On the other hand, the QMC calculations of Ref. 13 should be directly comparable to our data. The corresponding estimates of ϕ are in a rather good agreement with our experiments, as well as with those on the Br-substituted DTN compound.¹³ A relevant theoretical study of β is presently lacking. The claimed observation¹² of values consistent with $z = d$ can be due to overinterpretation of limited dataset. Evidence is mounting for the breakdown of the $z = d$ hypothesis in the Bose Glass to BEC quantum phase transition.

This work is partially supported by the Swiss National Fund under project 2-77060-11 and through Project 6 of MANEP. ML acknowledges support from DanScat. We thank Dr. V. Glazkov for his involvement in the early stages of this project.

* <http://www.neutron.ethz.ch/>

¹ T. Giamarchi, C. Ruegg, and O. Tchernyshev, Nature Physics **4**, 198 (2008).

² T. Giamarchi and H. J. Schulz, Europhys. Lett. **3**, 1287 (1987).

³ M. P. A. Fisher, P. B. Weichman, G. Grinstein, and D. S. Fisher, Phys. Rev. B **40**, 546 (1989).

⁴ O. Nohadani, S. Wessel, and S. Haas, Phys. Rev. Lett. **95**, 227201 (2005).

⁵ P. B. Weichman and R. Mukhopadhyay, Physical Review Letters **98**, 245701 (2007).

⁶ P. B. Weichman and R. Mukhopadhyay, Physical Review B **77**, 214516 (2008).

⁷ A. Priyadarshree, S. Chandrasekharan, J.-W. Lee, and

H. U. Baranger, Physical Review Letters **97**, 115703 (2006).

⁸ H. Meier and M. Wallin, arXiv:1109.3022v2.

⁹ H. Manaka, A. V. Kolomiets, and T. Goto, Phys. Rev. Lett. **101**, 077204 (2008).

¹⁰ H. Manaka, H. A. Katori, O. V. Kolomiets, and T. Goto, Phys. Rev. B **79**, 092401 (2009).

¹¹ T. Hong, A. Zheludev, H. Manaka, and L.-P. Regnault, Phys. Rev. B **81**, 060410 (2010).

¹² F. Yamada, H. Tanaka, T. Ono, and H. Nojiri, Phys. Rev. B **83**, 020409 (2011).

¹³ R. Yu, L. Yin, N. S. Sullivan, J. S. Xia, C. Huan, A. Paduan-Filho, N. F. Oliveira Jr., S. Haas, A. Steppke, C. F. Miclea, F. Weickert, R. Movshovich, E.-D. Mun, V. S.

- Zapf and T. Roscilde, arXiv:1109.4403v2.
- ¹⁴ E. Wulf, S. Mühlbauer, T. Yankova, and A. Zheludev, Phys. Rev. B **84**, 174414 (2011).
 - ¹⁵ M. B. Stone, I. Zaliznyak, D. H. Reich, and C. Broholm, Phys. Rev. B **64**, 144405 (2001).
 - ¹⁶ M. B. Stone, C. Broholm, D. H. Reich, O. Tchernyshyov, P. Vorderwisch, and N. Harrison, Phys. Rev. Lett. **96**, 257203 (2006).
 - ¹⁷ M. B. Stone, C. Broholm, D. H. Reich, P. Schiffer, O. Tchernyshyov, P. Vorderwisch, and N. Harrison, New Journal of Physics **9**, 31 (2007).
 - ¹⁸ M. B. Stone, I. A. Zaliznyak, T. Hong, C. L. Broholm, and D. H. Reich, Nature **440**, 187 (2006).
 - ¹⁹ T. Yankova, D. Hüvonen, S. Muehlbauer, D. Schmidiger, E. Wulf, S. Zhao, A. Zheludev, T. Hong, V. O. Garlea, R. Custelcean, G. Ehlers, arXiv:1110.6375v1.
 - ²⁰ G. Marcotrigiano, L. Menabue, and G. C. Pellacani, Inorganic Chemistry **15**, 2333 (1976).
 - ²¹ $C(T) = b - a_{l,r}\gamma^{-2} - a_{l,r}(|T - T_c|^{\alpha_{l,r}} + \gamma^2)^{-1}$ where γ is a broadening parameter, b is background, $a_{l,r}$ and $\alpha_{l,r}$ are scaling constants and scaling exponents on the left and right side of T_c respectively.
 - ²² F. Semadeni, B. Roessli, and P. Boni, Physica B **297**, 152 (2001).
 - ²³ M. Popovici, Acta Cryst. **A31**, 507 (1975).
 - ²⁴ A full account of the inelastic neutron scattering experiments are beyond the scope of the present paper that is focused on the field-induced transitions. It will be made elsewhere.
 - ²⁵ Lower limit on ϕ follows from Eqs. 5.8 and 5.10 in Ref. 3 as follows $T_c \propto [\rho_s(0)]^x$ and $\rho_s(0) \propto (\rho - \rho_c)^\zeta$ with $x = z/(d+z-2)$ and $\zeta = \nu(d+z-2)$. Since magnetization and boson density are proportional $(\rho - \rho_c) \propto (H - H_c)$ one obtains $T_c \propto (H - H_c)^{x\zeta} = (H - H_c)^{z\nu}$. Since $\nu \geq 2/d$ [Eq. 3.24 in Ref. 3] for all dimensions in the presence of disorder and $z = d$ we obtain dimension invariant $\phi \geq 2$.
 - ²⁶ R. Yu, S. Haas, and T. Roscilde, Europhys. Lett. **89**, 10009 (2010).